

Topological Interpretation of the Dirac Equation and Geometrical Foundations of the Gauge Invariance Idea

O.A. Ol'khov*

*Institute of Chemical Physics, Russian Academy of Sciences, 119991,
Moscow, Kosygin Str.,4, Russia*

Soon after the Yang-Mills work, the gauge invariance became one of the basic principles in the elementary particles theory. The gauge invariance idea is that Lagrangian has to be invariant not only with respect to the coordinates transformations corresponding to the Lorentz group (external symmetry). It is supposed that Lagrangian has also to be invariant with respect to wave functions (not coordinates) transformations corresponding to some additional groups (so-called "internal symmetry groups"). Useful though this idea is, there is no satisfactory understanding of the above additional symmetries origin, and the gauge invariance is considered as an auxiliary theoretical hypotheses. We propose a new, topological interpretation of the basic quantum mechanical equation — the Dirac equation, and within the framework of this interpretation the notions of internal symmetry and gauge invariance bear a simple geometrical meaning and are natural consequences of the basic principles of the proposed geometrical description. According to this interpretation the Dirac equation proves to be the group-theoretical relation that accounts for the symmetry properties of a specific 4-manifold — localized microscopic deviation of the space-time geometry from the Euclidean one. This manifold covering space plays the role of the internal space, and the covering space automorphism group plays the role of the gauge group.

1. Introduction

At high energies, the quantum mechanics formalism is now considered as a description of interacting quanta of different physical wave fields: electron–positron field, quark field, etc. At low energies, where the probability of new particles appearance is negligible, above formalism is considered as

a description of interacting pointlike particles having some wave properties (wave–corpuscular duality concept). In this presentation we show that the Dirac equation can be considered as a relationship between some geometry and group-theory notions. In result, this relationship can be considered as a description of symmetry proprieties of some specific microscopic space–time deformation. The macroscopic deformation of the space–time is the basic idea of a geometrical interpretation of the gravitational field within the framework of the general theory of relativity. So, the suggested interpretation, in some sense, is an attempt to extend the general relativity ideology to the description of microscopic quantum objects.

There is another aspect of this contribution. There were many attempts to find out some new interpretation of quantum mechanics formalism. On the other hand, in parallel, there were many attempts to geometrize electromagnetic and other physical fields. The distinctive feature of this work is that it combines both above problems within one idea. Namely, we show that the equations for the gauge electromagnetic field and the equations for this field sources can be considered as a description of a single geometrical object. Some preliminary results were published earlier [1-5].

This paper is organized as follows. In Section 2 we show that the Dirac equation for a free particle can be interpreted as a group-theoretical description of the specific space–time deformation, namely, the closed nonorientable four-dimensional manifold (4-manifold). Differentiation operators and Dirac's matrices in this equation prove to be some symmetry-group operators and the wave function proves to be a basis vector of the corresponding symmetry-group representation. In Section 3 we extend the above interpretation to the Dirac equation for a charged particle interacting with its own electromagnetic field. Within geometrical interpretation, this equation happens to be a group-theoretical relation written for some curved 4-space that is a standard geometrical characteristic of a closed 4-manifold (universal covering space of the manifold). This space proves to be the analogy of so called particle "internal"space — a notion additionally introduced into the relativistic

quantum mechanics. So, within the geometrical interpretation, the notions of internal space and its internal symmetry are not some additional attributes of the theory but are natural consequences of the geometrical description. Section 4 contains conclusion and final remarks. In Appendix, we illustrate by the simplest example (one-dimensional closed manifold) the possibility of the manifold description by means of linear differential equations.

2. Equation for free particle

Let us first consider the Dirac equation for a free particle [6],

$$i\gamma_\mu\partial_\mu\psi = m\psi, \quad (1)$$

where $\partial_\mu = \partial/\partial x_\mu$, $\mu = 1, 2, 3, 4$, γ_μ are the Dirac matrices, $\psi(x)$ is the Dirac bispinor, $x_1 = t, x_2 = x, x_3 = y$, and $x_4 = z$. There is a summation over repeating indices with a signature $(1, -1, -1, -1)$. Here, $\hbar = c = 1$. The solution to Eq.(1) has the form of a plane wave

$$\psi = u \exp(-ip_\mu x_\mu), \quad (2)$$

where the 4-momentum p_μ satisfies the relation

$$p_1^2 - p_2^2 - p_3^2 - p_4^2 = m^2. \quad (3)$$

The interpretation of Eq.(1) is, in fact, defined by the interpretation of its solution (2). The up-to-date interpretation depends on the particle energy. For $c \rightarrow \infty$ (nonrelativistic limit), $|\psi(x)|^2$ is interpreted as a probability to detect the particle at point x . The wave function $\psi(x)$ in (2) is said to represent, in this case, the "probability traveling wave" (the wave-corpuscular duality concept). For $v \sim c$, the probability of the new particles appearance needs to be taken into account. In this case, the wave function is interpreted as the annihilation and creation operator for the corresponding field quantum.

Let us show that the geometrical and symmetry characteristics of a certain closed space-time 4-manifold are encoded in Eqs.(1–3) by means of

the group-theory operators. It is now be helpful to rewrite Eqs.(1–3) in a more convenient form. We rewrite Eq.(1) as

$$(l_1^{-1}\gamma_1 T_1 - l_2^{-1}\gamma_2 T_2 - l_3^{-1}\gamma_3 T_3 - l_4^{-1}\gamma_4 T_4)\psi = l_m^{-1}\psi, \quad (4)$$

where operator T_μ ($\mu = 1, 2, 3, 4$) has the form

$$T_\mu = i(2\pi)^{-1}l_\mu\partial_\mu, \quad l_\mu = 2\pi p_\mu^{-1}, \quad l_m = 2\pi m^{-1}. \quad (5)$$

We rewrite the solution (2) as

$$\psi = u \exp(-2\pi i x_1 l_1^{-1} + 2\pi i x_2 l_2^{-1} + 2\pi i x_3 l_3^{-1} + 2\pi i x_4 l_4^{-1}). \quad (6)$$

We also rewrite the conservation law (3) as

$$l_1^{-2} - l_2^{-2} - l_3^{-2} - l_4^{-2} = l_m^{-2}. \quad (7)$$

Note that all quantities in Eqs (4–7) has the dimensionality of length.

Operators T_μ in Eq.(5) and functions $\psi(x)$ in Eq.(6) are related by the following equation.

$$\psi'(x'_\mu) \equiv T_\mu \psi(x'_\mu) = \psi(x_\mu), \quad x'_\mu = x_\mu + l_\mu. \quad (8)$$

Within a group theory the relation (8) means that T_μ is the operator representation of a discrete group of parallel translations along x_μ axis with the generator l_μ . The relation (8) means also that the function $\psi(x)$ is a basic vector of this representation [7].

In addition, being a four-component spinor, the function $\psi(x)$ is related to the four-row Dirac matrices γ_μ by the equations [8,9]

$$\psi'(x') = \gamma_\mu \psi(x), \quad (9)$$

where $x \equiv (x_1, x_2, x_3, x_4)$, $x' \equiv (x_1, -x_2, -x_3, -x_4)$ for $\mu = 1$, and $x' \equiv (-x_1, x_2, -x_3, -x_4)$ for $\mu = 2$, and so on. Within the group theory, Eqs.(9) mean that the matrices γ_μ are a matrix representation of the group of reflections along three axes perpendicular to the x_μ axis. Relation (9) means

also that the function $\psi(x)$ is a basic vector of the above symmetry-group representation.

A parallel translation with simultaneous reflection in the direction perpendicular to the translation is often spoken of as "sliding symmetry"[10]. Thus, we see that the operators

$$P_\mu = \gamma_\mu T_\mu. \quad (10)$$

form the representations of sliding symmetry group (sliding symmetry in the direction $0x_\mu$). Using the above notation, we can rewrite Eq.(1) as

$$l_\mu^{-1} P_\mu \psi = l_m^{-1} \psi, \quad (11)$$

where, as we established before, the function $\psi(x)$ can be interpreted as a basic vector of the above sliding symmetry group that operates in the four-dimensional space, in which Eqs.(1) and (11) are written.

There is an important moment of the presentation at this point. Eq.(1) is usually supposed to be written in the four-dimensional space-time that has no any sliding symmetry. The physical space-time has a symmetry only with respect to translations and 4-rotations (Poincaré group). This symmetry reflects the space-time homogeneity and, therefore, the Dirac equation (1) is an invariant with respect to this symmetry group [6]. What does the appearance of a new additional symmetry means? The suggestion about the reasons for an appearance, within mathematical formalism, of a new additional symmetry is a basis of the proposed geometrical interpretation of the Dirac equation (1).

Our main suggestion is that the Dirac equation (1) describes some special microscopic distortion of the physical space-time, and it is necessary to have some auxiliary four-dimensional space with the sliding symmetry to represent the distortion geometrical proprieties mathematically. The objects of this kind are known in geometry. These are so called closed topological manifolds. Such manifold is defined as any of the objects that can be obtained from some definite geometrical object by its deformation without discontinuities [10]. The above mentioned auxiliary space is spoken of as the manifold

universal covering space. The symmetry group of this space coincides with the so-called fundamental group of the closed manifold. This group is the group of different classes of closed paths starting and ending at the same point of the manifold and this group is one of the important topological characteristics of manifolds.

Thus, our suggestion is that the "discovered" sliding symmetries along four axes are symmetries of the four-dimensional universal covering space of the closed space-time manifold. It means that the fundamental group of this manifold is a group of four sliding symmetries. These four sliding symmetries correspond to four possible classes of closed paths. The Dirac equation (1) is supposed to be written for the manifold covering space, and Eq.(11) has to be considered as the group-theoretical relation that leads to the restriction (7) for the four possible values of l_μ (l_μ is a length of the closed path for one of the four possible classes).

By now, the topology of four-dimensional manifolds is understood to a much lesser extent than the topology of, e.g, two-dimensional manifolds. Only two-dimensional manifolds are classified in detail, and the parameters of their symmetry groups are known [10]. For this reason, we have no opportunity to identify our space-time 4-manifold with any well investigated topological object. So, we will attempt to invoke a possible analogy with some two-dimensional manifold having formally almost the same proprieties as the four-dimensional space-time manifold that we introduced before. The fundamental group of this two-dimensional manifold is generated by two sliding symmetries, and its universal covering space is a two-dimensional Euclidean space. We have in mind the so-called Klein bottle. The Klein bottle is one of the simplest examples of nonorientable geometrical objects and it can be obtained by gluing together two nonorientable Möbius strips [10]. So we can suggest that Eq.(11) describes microscopic closed nonorientable four-dimensional space-time manifold similar formally to the two-dimensional Klein bottle. Microscopic sizes of this manifold are defined by microscopic values of its length parameters l_μ and l_m . For example, $l_m = 2\pi\hbar/mc \sim 10^{-10}$

cm., if m is the electron mass.

3. Geometrical interpretation of the notions of "internal space", "internal symmetry", and "gauge invariance"

There is, of course, a question whether the suggested geometrical interpretation of Eg.(1) has any advantages in compare with the traditional one. We will show below that the geometrical interpretation resolves one of the problems of the relativistic quantum mechanics that has yet no solution within the framework of traditional interpretation.

One of the basic idea of the relativistic quantum mechanics is the suggestion that the basic equations have to be invariant with respect to some additional "internal" or "gauge" symmetry groups that differ from the "external" Poincaré group and that operate in some additional "internal space" of a particle. This suggestion is usefully employed in the gauge theory of elementary particles but there is no satisfactory explanation of these additional symmetries [11,12]. We will show below that, within the framework of the geometrical interpretation, the notions of internal symmetry, internal space, and gauge invariance bear a simple geometrical meaning and are natural consequences of the basic principles of geometrical description. We have seen in the previous section that the geometrical interpretation of Eq.(1) introduces into consideration the additional space—the universal covering space of 4-manifold. This fact will serve as a guide for geometrical interpretation of the "internal space."

We will consider the problem for a case of the classical Dirac electron–positron field interacting with its own gauge electromagnetic field. The system of equations for this case (Dirac and Maxwell equations) has the form [6]

$$i\gamma_\mu(\partial_\mu - ieA_\mu)\psi = m\psi. \quad (12)$$

$$F_{ik} = \partial_k A_i - \partial_i A_k, \quad (13)$$

$$\partial_i F_{ik} = j_k. \quad (14)$$

Here, e and m are electron charge and mass, respectively, A_l are 4-potentials of electromagnetic fields, and the current j_k is defined by the relation

$$j_k = e\psi^+\gamma_k\psi, \quad (15)$$

where $\psi^+ = \psi^*\gamma_1$.

The only distinction of Eq.(12) from Eq.(1) is the replacement of conventional derivatives ∂_μ by so-called long derivatives $(\partial_\mu - ieA_\mu)$. Inserting electromagnetic 4-potentials in Eq.(1) by means of the above-mentioned replacement is a consequence of the suggestion that Eq.(12) is the invariant with respect to the phase transformation of the wave function [11,12]

$$\psi \rightarrow \psi' = \psi \exp[ie\alpha(x)], \quad (16)$$

where $\alpha(x)$ is a parameter depending on the space-time coordinate x . This invariance (named as "gauge invariance") may be realized only by inserting electromagnetic 4-potentials into Eq.(1) as it is done in Eq.(12). In so doing, we obtain the following gauge relation for A_l [11,12].

$$A_l \rightarrow A'_l = A_l + \partial_l\alpha(x). \quad (17)$$

(Note that the above transformation leaves F_{ik} unchanged). Relation (16) does not directly include a space-time (external) coordinates transformation, and this fact is said to be the reflection of some "internal space" symmetry. As we mentioned before, the nature of this internal space and the origin of its gauge symmetry have no satisfactory explanation [11].

Let us now look at requirements of the gauge invariance (16,17) within the framework of the possible geometrical interpretation of Eq.(12). Within the geometrical approach, we have to consider Eq.(12) (similarly to Eq. (1)) as a group-theoretical relation for a four-dimensional (may be non-Euclidean) universal covering space of some space-time 4-manifold. But it will be the case only if we could prove that the long derivatives in Eq.(12) can be considered within a differential geometry as so-called covariant derivatives for some non-Euclidean space. Moreover, we have to find out transformations

of this space coordinates, that correspond to a gauge transformation of wave function (16), and we must prove that these coordinate transformations lead to the gauge relation (17).

It happened that the above-mentioned mathematical problem was long ago solved by Weil, who considered a different physical problem [15]. Weil has attempted to represent an electromagnetic field (analogously to a gravitational field) as a manifestation of the physical space-time curvature. He has proved, that the long derivatives in (12) $(\partial_l - ieA_l)$ can be, indeed, considered as covariant derivatives for the special curved 4-space (the space with semimetrical parallel translation). In this case, the values ieA_l play a role of the space specific geometrical characteristic—"connectivity". What is more, Weil has showed that gauge transformations (16,17) may be considered as a consequence of the following transformation of the space-time metric g_{ik} :

$$g'_{ik} = \lambda(x)g_{ik}. \quad (18)$$

The infinitesimal element of the 4-space length (the 4-interval ds) is defined by the relation [13]

$$ds^2 = g'_{ik}dx_i dx_k.$$

Therefore, transformation (18) corresponds to the 4-space isotropic stretching depending on space coordinates.

The real physical space-time has no any curvature due to an electromagnetic field. So, the "Weil space" was considered earlier just as a methodological result having nothing to do with the reality. But we see that, within the geometrical approach, the Weil space does not have to be considered as the curved real physical space-time. It has to be considered as a covering space of the 4-manifold whose proprieties are described by Eq.(12). This space (being a formal geometrical construction) plays the role of "internal"space, and gauge transformations (17,18) look now like natural manifestation of the Eq.(12) invariance about those covering space transformations that do not change its symmetry (Recall that the covering space symmetry defines all proprieties of the above-mentioned 4-manifold).

4. Conclusion and final remarks

The above considerations can be summarized as follows.

1. Equation (12), as well as Eq.(1), can be considered as a description of some microscopic closed space-time 4-manifold—specific microscopic localized deviation of the space geometry from the Euclidean one.

2. Gauge transformation of the wave function (16) corresponds to such coordinates transformation of the manifold covering space that does not change its symmetry and, therefore, does not change the mathematical description of the manifold. The manifold covering space proves to be non-Euclidean "Weil space."

3. Thus, the gauge invariance of Eq.(12) proves not to be some additional theoretical principle, but it happens to be a natural consequence of the basic principles of the proposed geometrical interpretation. The covering space plays the role of so called "particle internal space".

As we see, the next steps look as follows. First, it should be a geometrization of another known gauge groups such as SU(2)-group and SU(3)-group, and then it should be a geometrical interpretation of the secondary quantization procedure. The discussion of these problems will be presented elsewhere.

And the last remark. It is accepted that the absence of hidden parameters is the necessary condition for validity of any new quantum mechanics interpretation [16,17]. The suggested geometrical interpretation does not include any hidden parameters.

Appendix

To my knowledge, the topological manifolds were not before identified by the linear differential equations like Eq.(1). We will demonstrate, by a simplest example, how linear differential equations can describe the manifold topological and metric characteristics.

Let us consider the simplest closed connected manifold, namely, a one-dimensional manifold S^1 homeomorphic with (equivalent to) ring. It is equivalent

lent in the sense that it can be represented as any of the objects obtained from ring by its deformation without discontinuities. Different classes of closed paths starting and ending at the same point of the ring form its fundamental group and this group is the topological invariant of our manifold. The fundamental group of S^1 is the group of integers (Z group)[18]. In turn, Z group is isomorphous with the group of parallel translations along a straight line with one generator that we denotes as l , and this straight line (x axis) serves as universal covering space of our manifold (one-dimensional Euclidean space). So, the universal covering space of the ring is an one-dimensional Euclidean space with additional symmetry (the parallel translations group). The topological type of our manifold does not depend on the concrete value of the parameter l . This parameter is the manifold additional metric (not topological) characteristic that sets limits on the ring possible configurations. We will consider l as a given, preassigned quantity, $l = l_0$, and this quantity must be reflected along with the ring fundamental group within the manifold mathematical description (this restriction on l is an analogy of restriction (7) that appeared within the framework of the Dirac equation topological interpretation).

As we noticed in Section 2, the operator

$$T_x = \left(\frac{il_0}{2\pi} \right) \frac{d}{dx} \quad (A1)$$

can be considered as the operator representation of a discrete group of parallel translations along x axis with the generator l_0 . The basic vectors of this representation are represented by the functions $\varphi(x)$

$$\varphi(x) = \exp \left(-2\pi i \frac{x}{l_0} \right), \quad (A2)$$

because [7]

$$\varphi'(x') \equiv T_x \varphi(x') = \varphi(x), \quad x' = x + l_0. \quad (A3)$$

Thus, both the ring fundamental group and the metrical restriction on the length of the parameter of this group are defined by relationship (A3).

Thus, the manifold is fully identified by the linear differential equation

$$\left(\frac{il_0}{2\pi}\right) \frac{d\varphi}{dx} = \varphi(x + l_0) = \varphi(x),$$

or

$$i \frac{d\varphi}{dx} = m_0 \varphi, \quad m_0 = \frac{2\pi}{l_0}. \quad (A4)$$

Therefore, Eq.(A4) can be considered as one of the possible mathematical description of the ring topological and metrical properties. Of course, there may be another physical and geometrical interpretations of Eq.(A4).

*E-mail address: olega@gagarinclub.ru

1. O.A. Olkhov, in Proceedings of Int.Workshop on High Energy Physics and Field Theory, Protvino, Russia, June 2001, p.327.
2. O.A. Olkhov, in Proceedings of "Photon 2001", Ascona, Switzerland, September 2001, p.360.
3. O.A. Olkhov, in Proceedings of Int.Conf. on New Trends in High-Energy Physics, Yalta (Crimea), September 2001, p.324.
4. O.A. Olkhov in Proceedings of "Group 24", Paris, France, July 2002. p. 363.
5. O.A. Olkhov, Moscow Institute of Physics and Technology, Preprint No.2002-1, Moscow, 2002.
6. J.D. Bjorken and S.D. Drell, *Relativistic Quantum Mechanics*. (McGraw-Hill Book Company, 1964).
7. M. Hamermesh, *Group Theory and Its Application to Physical Problems*. (Argonne National Laboratory, 1964).
8. A.I. Achiezer, S.V. Peletminski, *Fields and Fundamental Interactions*. (Naukova Dumka, Kiev, 1986), Ch.1
9. E. Cartan, *Leçons sur la théorie des spineurs*. (Actualités scientifiques et industrielles, Paris, 1938).
10. See, e.g., H.S.M. Coxeter, *Introduction to Geometry*. (John Wiley and Sons, Inc., N.Y.-London, 1961).
11. G. Kane, *Modern Elementary Particle Physics*. (Addison-Wesley Publishing Company, Inc., 1987).

12. L.B. Okun, *Leptons and Quarks*. (North-Holland Publishing Co., 1982).
13. J.A. Schouten, *Tensor Analysis for Physicists*. (Clarendon Press, Oxford, 1951).
14. J.A. Schouten and D.J. Struik, *Einführung in die neue Methoden der Differentialgeometrie*. (Groningen-Batavia, Noordhoff, 1935), Vol.1.
15. H. Weyl, *Gravitation und Electrizzität*. (Preuss. Akad. Wiss., Berlin, 1918); Z.f.Phys. **56**, 330 (1929).
16. J.S. Bell, Rev.Mod.Phys. **38**, 447 (1966)
17. J.V. Neumann, *Mathematische grundlagen der quantenmechanik*. (Springer, Berlin, 1935).
18. See, e.g., B.A. Dubrovin, S.P. Novikov, A.T. Fomenko, *Modern Geometry. Methods and Applications*. (Nauka, Moscow, 1986).